

Methodological Challenges of Mixed Methods Intervention Evaluations

Hennie R. Boeije,¹ Sarah J. Drabble,² and Alicia O’Cathain²

¹Faculty of Social and Behavioural Sciences, Department of Methodology and Statistics, Utrecht University, The Netherlands, ²School of Health and Related Research, University of Sheffield, United Kingdom

Abstract. This paper addresses the methodological challenges that accompany the use of a combination of research methods to evaluate complex interventions. In evaluating complex interventions, the question about effectiveness is not the only question that needs to be answered. Of equal interest are questions about acceptability, feasibility, and implementation of the intervention and the evaluation study itself. Using qualitative research in conjunction with trials enables us to address this diversity of questions. The combination of methods results in a mixed methods intervention evaluation (MMIE). In this article we demonstrate the relevance of mixed methods evaluation studies and provide case studies from health care. Methodological challenges that need our attention are, among others, choosing appropriate designs for MMIEs, determining realistic expectations of both components, and assigning adequate resources to both components. Solving these methodological issues will improve our research designs and provide further insights into complex interventions.

Keywords: qualitative research, randomized controlled trials, mixed methods intervention evaluation, process evaluation

Randomized controlled trials (RCTs) are used to test whether interventions are effective. Increasingly researchers are using mixed methods intervention evaluations – which include an RCT – to address a range of questions related to understanding the effectiveness of an intervention (Creswell, Fetters, Plano Clark, & Morales, 2009; Drabble & O’Cathain, 2015). In this article we address the methodological challenges involved in undertaking mixed methods intervention evaluations. We describe the benefits of using a mixed methods evaluation that includes quantitative and qualitative methods rather than only an RCT. We focus on the field of health research in which RCTs are common and we have some expertise.

In the field of health research an RCT is defined as “...an experiment in which two or more interventions, possibly including a control intervention or no intervention, are compared by being randomly allocated to participants. In most trials one intervention is assigned to each individual but sometimes assignment is to defined groups of individuals (e.g., in a household) or interventions are assigned within individuals” (The Cochrane Collaboration, 2013, see definition of randomized controlled trial).

Interventions can be drugs, devices, or services in clinical settings, but can also be designed to impact on health behaviors in a range of settings such as schools and the

community. Examples of public health interventions are peer support delivered in schools to aid smoking cessation among young people (Campbell et al., 2008; Munro & Bloor, 2010) or improving healthy eating in children (Thomas et al., 2003).

An industry has grown up around undertaking RCTs of drugs. Yet much of what is evaluated in health is more complex than a drug and is known as a “complex intervention.” A complex intervention has many components or active ingredients (Campbell et al., 2007; Munro & Bloor, 2010; Oakley, Strange, Bonell, Allen, & Stephenson, 2006), which combine independently and interdependently (Campbell et al., 2007), making the whole more than the sum of its parts (Hawe, Shiell, & Riley, 2004; Oakley et al., 2006). The complexity of the intervention can take different forms such as the variation in behaviors of the people delivering or receiving the intervention, the different groups or organizations affected by the intervention, and the variation in outcomes (Medical Research Council, 2008). Examples of complex interventions include the introduction of lifestyle interventions for people with obesity, or the reorganization of the way in which a service is delivered to improve quality of care (Dixon-Woods, Bosk, Aveling, Goeschel, & Pronovost, 2011).

Within health research, the question “is an intervention – be it a drug, service or technology – effective in improving health?” is very important and RCTs are an excellent way of addressing the effectiveness question. But increasingly this question is expanded to understand how an intervention

is or is not effective and under what circumstances. This wider perspective requires methods that can address these issues, such as qualitative research (Britten, 2011; Pope & Mays, 1995). This is in line with arguments to move on from a focus on clinical interventions to encompass the arena of health care – stakeholders, for example, insurance, professionals, hospitals, government – and the everyday world in which patients experience health and health care (Barbour, 1999; Popay & Williams, 1998).

A mixed methods intervention evaluation includes an RCT and qualitative research. A number of studies have identified examples of this combination of RCT and qualitative research (Jansen, Foets, & de Bont, 2009; Lewin, Glenton, & Oxman, 2009). In a systematic review of international journal articles reporting qualitative research undertaken with trials in health, 296 published articles were identified between 2008 and September 2010 (O’Cathain, Thomas, Drabble, Rudolph, & Hewison, 2013). We draw on some of these examples, to explore methodological issues when undertaking mixed methods intervention evaluations.

Important Evaluation Questions

In this section we identify important questions relevant to evaluations of interventions which RCTs alone cannot address.

(1) *If the Intervention Was Effective in the Trial, Will it Be Effective in the Real World?*

Policy makers, clinicians, and patients want to implement evidence from RCTs in the real world. Implementing findings from interventions shown to be effective under experimental conditions requires information about which aspects of the intervention are essential for effectiveness (mechanisms of action) and the relevance of the context that the intervention was tested into other circumstances (transferability of evidence) (Dixon-Woods et al., 2011; Greenhalgh, Kristjansson, & Robinson, 2007; Newnham & Page, 2010; Pawson & Tilley, 2004). The need to address questions related to *how* and *why* interventions work through understanding mechanisms of action and the transferability of the evidence, has led to the inclusion of qualitative methods within these evaluative designs.

(2) *Why Was the Intervention not Effective?*

It can be time consuming and expensive to undertake RCTs. In theory, valuable evidence comes from RCTs which produce null results, that is, which show that an intervention was not effective. However, in practice, null RCTs can feel like a waste of resources unless researchers can understand why the intervention was not effective so that other researchers refrain from evaluating similar types of interventions and move toward those that have a more likelihood of being effective. Researchers use qualitative and quantitative methods to facilitate understanding of *why* interventions do not work by exploring whether

interventions have been delivered as planned or encountered barriers to successful implementation. For an example see Box 1.

Box 1: Case Study Breastfeeding In Groups (BIG) Study

Hoddinott, Britten, and Pill (2010) evaluated an intervention to assess how effective an intervention was to improve breastfeeding rates using an approach called “prospective mixed method embedded case studies” (p. 777). The breastfeeding in groups (BIG) trial was a cluster trial of 14 sites in Scotland in which 7 intervention sites increased group activities related to breastfeeding and 7 control localities retained their existing group activities. The research team utilized an ethnographic, realist evaluative methodology (Pawson & Tilley, 2004) to their trial design and conduct, which involved writing reflective diaries regarding meetings, telephone conversations, emails, and their thoughts and opinions about their experiences during the trial. They reflected on their backgrounds as a general practitioner and a former breastfeeding volunteer and how this affected their research roles. They hypothesized that the local environment in which the intervention took place would lead to differences between the intervention and control clusters. They conducted focus groups, interviews, and breastfeeding group observations during the study. Before analysing the trial outcomes, they utilized the qualitative research to devise an explanatory model of factors contributing to the success or failure of the sites’ ability to deliver the intervention. The trial itself had a null outcome, with a reduction in breastfeeding rates in three of the seven intervention sites. The model constructed from the qualitative data identified problems with leadership, less resources, and issues with the rooms where group sessions were held, which helped researchers to explain differences in outcomes between clusters.

(3) *What Is the Optimal Intervention to Test?*

It is important to understand an intervention so that large and expensive RCTs are not undertaken when it is not feasible for health professionals to deliver the intervention or the intervention is unacceptable to patients. For example, the intervention being tested in the RCT may be delivered in busy health service clinics which require health professionals to change their work practices. Qualitative methods such as interviews and focus groups with potential patients and those who will deliver the intervention can explore issues around feasibility and acceptability of interventions prior to an RCT, for example identifying how the intervention can be adapted to operate well in the context in which it will be tested and/or delivered in the real world. Thus an intervention and/or its

implementation can be optimized before undergoing expensive evaluation.

(4) *Will the RCT Work?*

RCTs can be challenging to run. For example, RCTs can struggle to recruit enough participants leading to low statistical power, or they can only recruit a narrow profile of the population that will actually use the intervention in the real world so they lack external validity (Bird, Arthur, & Cox, 2011). Questions about how to improve the conduct of the RCT can be important, and qualitative research has addressed this issue. For example, research has identified misunderstandings that lead to nonparticipation in the trial by interviewing health professionals recruiting patients for RCTs and patients approached for participation (Dixon-Woods, Jackson, Windridge, & Kenyon, 2006; Donovan et al., 2002). Observations of recruitment practices can identify reasons for misunderstanding such as poor communication.

(5) *Have We Really Understood the Complexities of What We Are Researching?*

The multifaceted nature and dependence on social context make complex interventions particularly challenging and costly to evaluate. The nature of complex interventions creates methodological challenges for the RCT (Campbell et al., 2007; Oakley et al., 2006) such as difficulties in standardizing the design and delivery of the intervention and understanding aspects of the local context in which the intervention is delivered (Medical Research Council, 2008). While an RCT may remain the most rigorous way to evaluate the effectiveness of an intervention, there is increasing acknowledgment of the contribution that qualitative methods can make to understand the complexity of interventions (Glenton, Lewin, & Scheel, 2011).

Methodological Challenges of Mixed Methods Intervention Evaluations

Although the indications for using a mixed methods intervention evaluation might be clear to researchers, it might be difficult to undertake one in practice. Lewin et al. (2009) examined 30 trials of complex interventions designed to change the organization of care where qualitative research was also used. In a systematic mapping review, O’Cathain et al. (2013) identified 296 articles published between January 2008 and September 2010 that reported the qualitative research undertaken with an RCT and mapped the focus of the qualitative research in relation to the RCT in practice. Based on these outcomes and a study of interviews with researchers undertaking mixed methods intervention evaluations (O’Cathain et al., 2014), we formulate methodological challenges in undertaking and reporting mixed methods evaluations.

Choosing an Appropriate Design

A common approach to describing the use of qualitative research with RCTs is a temporal framework, which considers how qualitative research can be used before, during, or after a trial (Creswell et al., 2009; Jansen et al., 2009; Lewin et al., 2009; Sandelowski, 1996). What design is appropriate depends on the questions that need to be addressed with both strands in the mixed methods evaluation design (Boeije, Slagt, & van Wesel, 2013). In a before-trial design qualitative data is collected and analyzed before the trial with the purpose of improving it. The focus of the qualitative research can be on the intervention, the trial conduct, or both. For an example, see Box 2.

Box 2: Adherence to Treatment of Cystic Fibrosis in Adolescents

Marciel et al. (2010) used mixed methods (focus groups, interviews, survey) before an RCT in the United States to adapt a peer support group intervention for adolescents, which had previously been tested on preschool and school-age children. The intervention aimed to improve adherence to treatment for cystic fibrosis. Focus groups were conducted with healthcare professionals and interviews undertaken with adolescent patients, adult patients, and parents. The qualitative research identified that feelings of invincibility and lack of knowledge about the consequences of non-adherence made adherence in this age group particularly difficult. It also identified unanticipated benefits of the mobile phone technology used in the intervention which allowed adolescents to have direct contact with their healthcare team. In the interviews, parents identified time constraints, lack of motivation, and forgetting treatments as barriers to facilitating adherence. Interviews with adolescents confirmed the acceptability of the proposed intervention to them which facilitated confidence in the proposed intervention prior to testing in an RCT.

A during-trial design involves the collection of qualitative data during the definitive trial to understand how the intervention is implemented in practice. A typical during-trial design is the process-outcome evaluation where the process evaluation aims to explain the trial results by understanding how the intervention was implemented in practice. See Box 1 for an example. Researchers have argued that process evaluations should be an integral part of trials of complex interventions, exploring why and how interventions work or do not work (e.g., Linnan & Steckler, 2002; Munro & Bloor, 2010; Oakley et al., 2006; Siu, Shek, & Poon, 2009). A particular strength of process evaluations is the ability to distinguish between what is planned in a particular setting and what is actually done in practice (Aro, Smith, & Dekker, 2008). Specifically, process evaluations can distinguish between “interventions that are inherently faulty (failure of intervention concept or theory) and those that are badly delivered (implementation failure)”

(Oakley et al., 2006, p. 413). This can help to avoid what has been termed “type III errors” (Audrey, Holliday, Parry-Langdon, & Campbell, 2006), that is, where a trial has a null result due to inadequate implementation of the intervention rather than a failure in the intervention design itself. Process evaluations may be particularly useful in cluster or multisite trials to understand how the context influences the same intervention being delivered at different sites (Oakley et al., 2006) and when an intervention attempts to change the organization of healthcare delivery (Glenton et al., 2011). The UK MRC has recently issued guidance to help researchers design and implement good process evaluations (Moore et al., 2014).

An after-trial design involves collecting qualitative data after the trial has ended. Qualitative data collected after the trial can explore the longer term implementation of an intervention [Medical Research Council (MRC), 2000, 2008] or may help researchers to explain the results of the trial if process evaluation data collected during the trial are unable to achieve this. When the trial is over the question remains how the trial findings can be used in daily practice. Qualitative research can help to translate the outcomes in the real world by studying relevant outcomes from the participants’ point of view, studying how new practices are implemented, and how results are disseminated in the media and translated into policy and practice.

Being Clear About the Expectations for the Qualitative Research

There may be tension around expectations for the qualitative research and what can be delivered in practice. Although the qualitative research may enrich understanding of the trial findings, this understanding “will always be nuanced and qualified and rarely determinate” (Munro & Bloor, 2010, p. 710) and researchers must always remain open to the possibility of other interpretations of the data. Expectations also pertain to the resources that are spent on the qualitative component: no unreasonable limits should be placed on the time, skill, and money required to complete an in-depth analysis which has quality and credibility (O’Cathain et al., 2014).

When it is clear what qualitative research should achieve, it should be part of the research protocol. It is standard practice to publish a protocol for an RCT so that researchers can be held to account, for example in their choice of the primary outcome. Oakley et al. (2006) argue that process evaluations should prospectively specify a set of research questions and identify the processes to be studied, the methods to be used, and the procedures for integrating the findings of the process evaluation with the results of the trial. Such a practice encourages planning of the qualitative research and sharing of this plan with the whole team of researchers. In turn, this can facilitate integration between relevant components of the study. Process evaluations can be published within trial protocols (e.g., Murphy

et al., 2010) or as a stand-alone process evaluation protocol (e.g., Ellard, Taylor, Parsons, & Thorogood, 2011; Grant, Dreischulte, Treweek, & Guthrie, 2012). Guidance has been produced to help researchers write proposals for using qualitative research with RCTs (Drabble, O’Cathain, Thomas, Rudolph, & Hewison, 2014).

Clearly Reporting How the Qualitative Research Has Enhanced the RCT

A key value of mixed methods research is that the whole is more than the sum of its parts (Barbour, 1999). For this to happen, one method must influence in some way the objectives, sampling, data collection, analysis, or interpretation of the other method within the study (O’Cathain, Murphy, & Nicholl, 2007; Small, 2011). Within mixed methods intervention evaluations, the expectation is that the qualitative research will work to enhance the trial (O’Cathain et al., 2013; Popay & Williams, 1998; Song, Sandelowski, & Happ, 2010). An example was given in Box 1, in which the integration of qualitative research explained how differences in staff attitudes and resources between clusters in a cluster RCT influenced differences in the primary outcome of breastfeeding rates between those clusters.

However, overview studies show that publications from mixed methods intervention evaluations often had no evidence of integration of the findings from the qualitative research and the trial (Lewin et al., 2009; O’Cathain et al., 2013). A challenge to all researchers engaged in mixed methods intervention evaluations is to explicitly report in journal articles the “yield” or insights gained from undertaking qualitative research and RCTs within the same study (Boeije et al., 2013; O’Cathain et al., 2007).

Integration may be difficult because some evaluation team members may not feel able to act on findings generated from the qualitative research. This might be an issue in a before-trial design where the findings may not be available at the time they are needed, the findings may not have credibility among key team members such as the lead researcher, the trialists may be unwilling to deviate from a particular path for their intervention and trial it, or the qualitative research may challenge established ways of thinking and practice. For example, Jansen et al. (2009) raised a concern that researchers actually focus on adapting the context to fit the intervention rather than considering how an intervention can be adapted to fit the context in which it will be delivered. Qualitative research that challenges such a mindset by suggesting a need to adapt the intervention to fit the context may be unacceptable to the wider team. To facilitate integration, the whole evaluation team may need to embrace a reflexive approach associated with qualitative research (Hesse-Biber, 2012). This happens best at an early stage of the study when teams can discuss how different values and beliefs shape their actions and reflect on their openness to different possibilities.

When a mixed methods intervention evaluation is complete, qualitative researchers and trialists may publish their findings in separate journals, each basing their articles solely on the findings from their part of the study. It is important that the timing of publication of articles allows for reporting of why the intervention was effective or not in the article reporting the trial.

Trial Governance

High-quality RCTs are governed by external trial steering committees and data monitoring committees, are conducted according to standardized operating procedures, and are reported in accordance with the CONSORT (Consolidated Standards of Reporting Trials) statement (Schulz, Altman, Moher, & for the CONSORT Group, 2010). This requires careful thought of the pros and cons of including the qualitative research in these practices and procedures (Cooper et al., 2014; Rapport et al., 2013). Qualitative research should not be blindly subsumed within all these procedures with an accompanying loss of its key strengths – its flexibility –, nor blindly excluded. An example of the careful thought needed is that discussion about the whole study at trial steering committees, rather than only the trial, can ensure more opportunities for integration (Rapport et al., 2013). Yet when to reveal findings from the qualitative research to the wider team may need careful thought if negative findings have the potential to demoralize the intervention deliverers or staff delivering the trial prior to the end of the trial (Cooper et al., 2014).

Conclusions

Increasingly when researchers evaluate interventions they are likely to use mixed methods intervention evaluations where they combine qualitative research with an RCT. This is a positive move driven by researchers understanding the complexity of the interventions they evaluate, the trials they conduct, and the environments in which they research. While this has become part of routine practice in some research communities, we have identified a number of methodological challenges with this approach in health research. None of the challenges presented are insurmountable and indeed reflecting on these challenges is likely to help research communities to understand how to maximize the benefits of mixed methods intervention evaluations.

Acknowledgments

This paper is based on a book chapter by Drabble and O’Cathain (2015), moving from randomized controlled

trials to mixed methods intervention evaluations. In S. Hesse-Biber and B. Johnson (Eds.), *The Oxford Handbook of Multimethod and Mixed Methods Research Inquiry*. Oxford University Press (ISBN: 9780199933624). It is also an adapted version of the keynote lecture by Alicia O’Cathain at the VI Conference of the European Association of Methodology held on July 24, 2014, Utrecht, The Netherlands.

References

- Aro, A. R., Smith, J., & Dekker, J. (2008). Contextual evidence in clinical medicine and health promotion. *European Journal of Public Health, 18*, 548–549.
- Audrey, S., Holliday, J., Parry-Langdon, N., & Campbell, R. (2006). Meeting the challenges of implementing process evaluation within randomized controlled trials: The example of ASSIST (A Stop Smoking in Schools Trial). *Health Education Research, Theory and Practice, 21*, 366–377.
- Barbour, R. S. (1999). The case for combining qualitative and quantitative methods in health services research. *Journal of Health Services Research and Policy, 4*, 39–43.
- Bird, L., Arthur, A., & Cox, K. (2011). “Did the trial kill the intervention?” Experiences from the development, implementation and evaluation of a complex intervention. *BMC Medical Research Methodology, 11*, 24. doi: 10.1186/1471-2288-11-24
- Boeije, H., Slagt, M., & van Wesel, F. (2013). The contribution of mixed methods research to the field of childhood trauma: A narrative review focused on data integration. *Journal of Mixed Methods Research, 7*, 347–369. doi: 10.1177/1558689813482756
- Britten, N. (2011). Qualitative research on health communication: What can it contribute? *Patient Education and Counseling, 82*, 384–388. doi: 10.1016/j.pec.2010.12.021
- Campbell, N. C., Murray, E., Darbyshire, J., Emery, J., Farmer, A., Griffiths, F., . . . Kinmonth, A. L. (2007). Designing and evaluating complex interventions to improve health care. *British Medical Journal, 334*, 455–459. doi: 10.1136/bmj.39108.379965
- Campbell, R., Starkey, F., Holliday, J., Audrey, S., Bloor, M., Parry-Langdon, N., . . . Moore, L. (2008). An informal school-based peer-led intervention for smoking prevention in adolescence (ASSIST): A cluster randomised trial. *Lancet, 371*, 1595–1602.
- Cooper, C., O’Cathain, A., Hind, D., Adamson, J., Lawton, J., & Baird, W. (2014). Conducting qualitative research within Clinical Trial Units: Avoiding potential pitfalls. *Contemporary Clinical Trials, 38*, 338–343.
- Creswell, J. W., Fetters, M. D., Plano Clark, V. L., & Morales, A. (2009). Mixed methods intervention trials. In S. Andrew & E. J. Halcomb (Eds.), *Mixed methods research for nursing and the health sciences* (pp. 161–180). Oxford, UK: Wiley-Blackwell.
- Dixon-Woods, M., Bosk, C. L., Aveling, E. L., Goeschel, C. A., & Pronovost, P. J. (2011). Explaining Michigan: Developing an ex post theory of a quality improvement program. *The Milbank Quarterly, 89*, 167–205.
- Dixon-Woods, M., Jackson, C., Windridge, K. C., & Kenyon, S. (2006). Receiving a summary of the results of a trial: Qualitative study of participants’ views. *British Medical Journal, 332*, 206–210. doi: 10.1136/bmj.38675.677963.3A

- Donovan, J., Mills, N., Smith, M., Brindle, L., Jacoby, A., Peters, T., ... Hamdy, F. (2002). Quality improvement report – Improving design and conduct of randomised trials by embedding them in qualitative research: ProtecT (PROstate TEsting for Cancer and Treatment) study. *British Medical Journal*, *325*, 766–769.
- Drabble, S. J., & O’Cathain, A. (2015). Moving from randomised controlled trials to mixed methods intervention evaluations. In S. Hesse-Biber & B. Johnson (Eds.), *The Oxford handbook of multimethod and mixed methods research inquiry* (pp. 406–425). Oxford, UK: Oxford University Press.
- Drabble, S. J., O’Cathain, A., Thomas, K. J., Rudolph, A., & Hewison, J. (2014). Describing qualitative research undertaken with randomised controlled trials in grant proposals: A documentary analysis. *BMC Medical Research Methodology*, *14*, 24.
- Ellard, D. R., Taylor, S. J. C., Parsons, S., & Thorogood, M. (2011). The OPERA trial: A protocol for the process evaluation of a randomised trial of an exercise intervention for Older People in Residential and Nursing Accommodation. *Trials*, *12*, 28. doi: 10.1186/1745-6215-12-28
- Glenton, C., Lewin, S., & Scheel, I. B. (2011). Still too little qualitative research to shed light on results from reviews of effectiveness trials: A case study of a Cochrane review on the use of lay health workers. *Implementation Science*, *6*, 53–57. doi: 10.1186/1748-5908-6-53
- Grant, A., Dreischulte, T., Treweek, S., & Guthrie, B. (2012). Study protocol of a mixed-methods evaluation of a cluster randomized trial to improve the safety of NSAID and antiplatelet prescribing: Data-driven quality improvement in primary care. *Trials*, *13*, 154–160.
- Greenhalgh, T., Kristjansson, E., & Robinson, V. (2007). Realist review to understand the efficacy of school feeding programmes. *British Medical Journal*, *335*, 858–861.
- Hawe, P., Shiell, A., & Riley, T. (2004). Complex interventions: How “out of control” can a randomised controlled trial be? *British Medical Journal*, *328*, 1561–1563.
- Hesse-Biber, S. (2012). Weaving a multimethodology and mixed methods praxis into randomised controlled trials to enhance credibility. *Qualitative Inquiry*, *18*, 876–889. doi: 10.1177/1077800412456964
- Hoddinott, P., Britten, J., & Pill, R. (2010). Why do interventions work in some places and not others: A breastfeeding support group trial. *Social Science and Medicine*, *70*, 769–778. doi: 10.1016/j.socscimed.2009.10.067
- Jansen, Y. J. F. M., Foets, M. M. E., & de Bont, A. (2009). The contribution of qualitative research to the development of tailor-made community-based interventions in primary care: A review. *European Journal of Public Health*, *20*, 220–226.
- Lewin, S., Glenton, C., & Oxman, A. D. (2009). Use of qualitative methods alongside randomised controlled trials of complex healthcare interventions: Methodological study. *British Medical Journal*, *339*, b3496. doi: 10.1136/bmj.b3496
- Linnan, L., & Steckler, A. (2002). Process evaluations for public health interventions and research. In A. Steckler & L. Linnan (Eds.), *Process evaluations for public health interventions and research* (pp. 1–23). San Francisco, CA: Jossey-Bass.
- Marciel, K. K., Saiman, L., Quittell, L. M., Dawkins, K., & Quittner, A. L. (2010). Cell phone intervention to improve adherence: Cystic fibrosis care team, patient, and parent perspectives. *Pediatric Pulmonology*, *45*, 157–164. doi: 10.1002/ppul.21164
- Medical Research Council. (2000). *A framework for the development and evaluation of RCTs for complex interventions to improve health*. London, UK: MRC. Retrieved from <http://www.mrc.ac.uk/Utilities/Documentrecord/index.htm?d=MRC003372>
- Medical Research Council. (2008). *Developing and evaluating complex interventions: New guidance*. London, UK: MRC. Retrieved from <http://www.mrc.ac.uk/Utilities/Documentrecord/index.htm?d=MRC004871>
- Moore, G., Audrey, S., Barker, M., Bond, L., Bonell, C., Hardeman, W., ... Baird, J. (2014). *Process evaluation of complex interventions*. UK Medical Research Council (MRC) guidance. Prepared on behalf of the MRC Population Health Science Research Network.
- Munro, A., & Bloor, M. (2010). Process evaluation: The new miracle ingredient in public health research? *Qualitative Research*, *10*, 699–713. doi: 10.1177/1468794110380522
- Murphy, S., Raisanen, L., Moore, G., Tudor Edwards, R., Linck, P., Williams, N., ... Moore, L. (2010). A pragmatic randomised controlled trial of the Welsh national exercise referral scheme: Protocol for trial and integrated economic and process evaluation. *BMC Public Health*, *10*, 352. doi: 10.1186/1471-2458-10-352
- Newnham, E. A., & Page, A. C. (2010). Bridging the gap between best evidence and best practice in mental health. *Clinical Psychology Review*, *30*, 127–142. doi: 10.1016/j.cpr.2009.10.004
- Oakley, A., Strange, V., Bonell, C., Allen, E., & Stephenson, J. (2006). Process evaluation in randomised controlled trials of complex interventions. *British Medical Journal*, *332*, 413–416.
- O’Cathain, A., Murphy, E., & Nicholl, J. (2007). Integration and publication as indicators of “yield” from mixed methods studies. *Journal of Mixed Methods Research*, *1*, 147–163. doi: 10.1177/1558689806299094
- O’Cathain, A., Goode, J., Drabble, S. J., Thomas, K. J., Rudolph, A., & Hewison, J. (2014). Getting added value from using qualitative research with randomised controlled trials: A qualitative interview study. *Trials*, *15*, 215.
- O’Cathain, A., Thomas, K. J., Drabble, S. J., Rudolph, A., & Hewison, J. (2013). What can qualitative research do for randomised controlled trials? A systematic mapping review. *BMJ Open*, *3*, e002889. doi: 10.1136/bmjopen-2013-002889
- Pawson, R., & Tilley, N. (2004). *Realist evaluation*. London, UK: Sage.
- Popay, J., & Williams, G. (1998). Qualitative research and evidence-based healthcare. *Journal of the Royal Society of Medicine*, *91*(Suppl. 35), 32–37.
- Pope, C., & Mays, N. (1995). Reaching the parts other methods cannot reach: An introduction to qualitative methods in health and health services research. *British Medical Journal*, *311*, 42–45.
- Rapport, F., Storey, M., Porter, A., Snooks, H., Jones, K., Peconi, J., ... Russell, I. (2013). Qualitative research within trials: Developing a standard operating procedure for a clinical trials unit. *Trials*, *14*, 54. doi: 10.1186/1745-6215-14-54
- Sandelowski, M. (1996). Using qualitative methods in intervention studies. *Research in Nursing and Health*, *19*, 359–364.
- Schulz, K. F., Altman, D. G., & Moher, D., for the CONSORT Group. (2010). CONSORT 2010 Statement: Updated guidelines for reporting parallel group randomised trials. *British Medical Journal*, *340*, c332. doi: 10.1136/bmj.c332
- Siu, A. M. H., Shek, D. T. L., & Poon, P. K. K. (2009). Evidence-based research in community rehabilitation: Design issues and strategies. *Hong Kong Journal of Occupational Therapy*, *19*, 20–26.

- Small, M. L. (2011). How to conduct a mixed methods study: Recent trends in a rapidly growing literature. *Annual Review Sociology, 37*, 57–86.
- Song, M., Sandelowski, M., & Happ, M. B. (2010). Current practices and emerging trends in conducting mixed methods intervention studies in the health sciences. In A. Tashakkori & C. Teddlie (Eds.), *Sage handbook of mixed methods research in social and behavioural research* (2nd ed., pp. 725–747). Thousand Oaks, CA: Sage.
- The Cochrane Collaboration. (2013). *Glossary*. Retrieved from <http://www.cochrane.org/glossary/5#letter>
- Thomas, J., Harden, A., Oakley, A., Oliver, S., Sutcliffe, K., Rees, R., . . . Kavanagh, J. (2003). *Children and healthy eating: A systematic review of barriers and facilitators*. London, UK: EPPI-Centre, Social Science Social Inequalities, Research Unit, Institute of Education, University of London.

Published online December 23, 2015

Hennie R. Boeije

Faculty of Social and Behavioural Sciences
Department of Methodology and Statistics
Utrecht University
P.O. Box 80140
3508 TC Utrecht
Tel. +31 30 2537761
E-mail h.boeije@uu.nl
